

References

- 1 Stoop JW. Pioneers in pediatric medicine: Willem Karel Dicke. *Eur J Pediatr* 150; 751: 1991.
- 2 Stheeman HA. De intestinale infantilismus en zijn behandeling met Ventraemon. *Ned Tijdschr Geneesk* 76; 4823: 1932.
- 3 Booth CC. History of coeliac disease. *Br Med J* 298; 527: 1989.
- 4 Dicke WK. Coeliac disease. Investigation of the harmful effects of certain types of cereal on patients with coeliac disease. Thesis University of Utrecht, 1950 (in Dutch). Second edition 1992 (in English).
- 5 Dicke WK. Simple dietary treatment for the syndrome of Gee-Herter. *Ned Tijdschr Geneesk* 85; 1715: 1941 (in Dutch).
- 6 Haas SV. Value of banana treatment in celiac disease. *Am. J. Dis. Child.* 1924;28:240.
- 7 Haas SV. Celiac disease, its specific treatment and cure without nutritional relapse. *JAMA* 1932;99:448.
- 8 Van De Kamer JH, ten Bokkel Huinink H and Weyers HA. Rapid method for the determination of fat in feces. *J Biol Chem* 177; 347: 1949.
- 9 Weyers HA. Fat absorption in normal and diseased neonates and children, especially in patients with coeliac disease. Thesis, University of Utrecht, The Netherlands, 1950 (in Dutch).
- 10 Van De Kamer JH. Coeliac disease, sprue syndrome and the metabolism of fat. A review of papers by Dicke, Frazer, Van De Kamer, Tegelaers, Trap and Weyers. *Voeding* 13; 1: 1952.
- 11 Van De Kamer JH, Weyers HA, Dicke KW. Coeliac disease IV. An investigation into the injurious constituents of wheat in connection with their action on patients with coeliac disease. *Acta Paed.* 42; 223 1953.
- 12 Anderson CM, Frazer AC, French JM, Gerrard JW, Simmons HG, Smellie JM. Coeliac disease: gastro-intestinal studies and the effects of dietary wheat flour. *Lancet* i; 836: 1952.
- 13 Söderhjelm AL. Dietbehandlingen vid coeliaki. *Nordisk Medicin* 10; 479: 1952
- 14 Krainick HG, Debatin F. Der schädliche Mehleffekt bei der kindlichen Cöliakie. *Klin Wschr* 31; 958: 1953.
- 15 Green PA, Wollaeger EE, Scudamore HH. Non-tropical sprue. Functional efficacy of small intestine after prolonged use of gluten-free diet. *JAMA* 171; 2157: 1959.
- 16 Peña A.S. History of coeliac disease. Dicke and the origin of the gluten-free diet. In: *Coeliac Disease, 40 years gluten-free*. Edited by: M.L. Mearin and C.J.J. Mulder.
- 17 Van Berge-Henegouwen GP, Mulder C.J.J. History of the discovery of the harmful effect of wheat products in celiac disease by Dr. Willem Karel Dicke: 50 years of gluten-free diet. 1993: in *Slesinger Gastroenterology*.

POSTULATIONS

- I If wheat and rye did not exist, is it possible that coeliac disease would not develop.
- II In the treatment of coeliac disease, rice flour, cornflour and potatoes can virtually be included in the diet from the outset.
- III The experience that anaemia is harmful to the organism, broadens the range of indications for tracheotomy.
- IV In a stubborn case of tendinitis of the m. supraspinatus tendon, one should consider resection of the acromion.
- V Infant and toddler clinics should become more involved in the prevention of infectious diseases.
- VI When treating burns, one should pay more attention to new ideas about how to prevent secondary infection.
- VII In connection with the development of chemo- and antibiotic therapy, it is not financially or economically responsible to extend sanatoria unless this is in the form of semipermanent buildings.
- VIII Group composition is very relevant to the results of a scientific study.
- IX In cases of compression symptoms of the trachea at a young age, after tuberculosis one considers anomalies of the great vessels as the possible cause.
- X It is to be recommended that the name scarlet fever be reserved for the exanthematous illness caused by haemolytic streptococci.