

**Chemistry 12**  
 June 2000 Provincial Examination  
**ANSWER KEY / SCORING GUIDE**

---

**CURRICULUM:**

<b>Organizers</b>	<b>Sub-Organizers</b>
1. Reaction Kinetics	A, B, C
2. Dynamic Equilibrium	D, E, F
3. Solubility Equilibria	G, H, I
4. Acids, Bases, and Salts	J, K, L, M, N, O, P, Q, R
5. Oxidation – Reduction	S, T, U, V, W

**Part A: Multiple Choice**

<b>Q</b>	<b>K</b>	<b>C</b>	<b>CO</b>	<b>PLO</b>	<b>Q</b>	<b>K</b>	<b>C</b>	<b>CO</b>	<b>PLO</b>
1.	A	K	1	A1	25.	A	U	4	K9
2.	D	K	1	A2	26.	B	U	4	L4, N3
3.	C	U	1	A3	27.	D	K	4	L6
4.	C	H	1	A5	28.	C	U	4	L11
5.	B	K	1	B5	29.	B	K	4	M1
6.	B	U	1	C5	30.	C	K	4	M2
7.	B	U	2	D6	31.	A	U	4	M4
8.	C	K	2	D9	32.	A	K	4	O2
9.	B	U	2	E2	33.	D	U	4	O4
10.	D	K	2	E2, 4	34.	B	U	4	P3
11.	C	K	2	F2	35.	C	U	4	P5
12.	C	H	2	F4	36.	A	U	4	P6
13.	A	U	2	F6	37.	B	H	4	Q5
14.	D	U	3	G5	38.	D	K	5	S1
15.	D	K	3	G8	39.	B	K	5	S2
16.	A	U	3	H2	40.	A	K	5	S2
17.	A	U	3	H4	41.	C	U	5	S3
18.	B	K	3	I2	42.	A	U	5	S6
19.	C	H	3	H2, I4	43.	C	K	5	T1
20.	D	H	3	I5	44.	D	U	5	T5
21.	D	K	4	J11	45.	B	H	5	U7
22.	C	K	4	K7	46.	D	U	5	U11
23.	D	U	4	J8	47.	B	U	5	V4
24.	A	U	4	K8	48.	A	U	5	W2

**Multiple Choice = 48 marks**

**Part B: Written Response**

<b>Q</b>	<b>B</b>	<b>C</b>	<b>S</b>	<b>CO</b>	<b>PLO</b>
1.	1	U	3	1	C2, C5
2.	2	U	3	2	B6, E2
3.	3	U	3	2	F6
4.	4	U	3	3	I5
5.	5	H	3	3	E2, H5, K5
6.	6	U	4	4	M5
7.	7	U	5	4	N1, N2, N3
8.	8	U	4	5	S4, U7
9.	9	U	2	5	U2, U11
10.	10	U	2	5	W8

**Written Response = 32 marks**

Multiple Choice = 48 (48 questions)

Written Response = 32 (10 questions)

**EXAMINATION TOTAL = 80 marks**

**LEGEND:**

**Q** = Question Number

**K** = Keyed Response

**C** = Cognitive Level

**B** = Score Box Number

**S** = Score

**CO** = Curriculum Organizer

**PLO** = Prescribed Learning Outcome

## PART B: WRITTEN RESPONSE

Value: 32 marks

Suggested Time: 50 minutes

**INSTRUCTIONS:** You will be expected to communicate your knowledge and understanding of chemical principles in a clear and logical manner.

Your steps and assumptions leading to a solution must be written in the spaces below the questions.

Answers must include units where appropriate and be given to the correct number of significant figures.

**For questions involving calculation, full marks will NOT be given for providing only an answer.**

1. a) Complete the steps in the following mechanism.

(1½ marks)

**Solution:**

*For Example:*

Step 1	$\text{NO} + \text{Pt} \rightarrow \underline{\text{NOPt}}$	← ½ mark
Step 2	$\text{NOPt} + \text{NO} \rightarrow \underline{\text{N}_2} + \underline{\text{O}_2\text{Pt}}$	← 1 mark
Step 3	$\text{O}_2\text{Pt} \rightarrow \text{O}_2 + \text{Pt}$	
Overall	$2\text{NO} \rightarrow \text{N}_2 + \text{O}_2$	

b) Define the term *reaction intermediate* and give an example from the completed mechanism above.

(1½ marks)

**Solution:**

*For Example:*


Definition: A substance which forms in one step of a mechanism and is used up in a later step.

} ← 1 mark

Example: NOPt or O<sub>2</sub>Pt

← ½ mark

2. Consider the observations for the following equilibrium:




Trial	Temperature °C	Colour
I.	10	light brown
II.	50	dark brown

a) Sketch the potential energy curve on the graph below for this equilibrium.

(1 mark)

**Solution:**



b) Explain the colour change using Le Châtelier's Principle.

(1 mark)

**Solution:**

*For Example:*

An increase in temperature causes the reaction to shift to the right and the  $[\text{NO}_2]$  increases.

← 1 mark

c) Other than changing temperature, what could be done to cause a shift to the left? (1 mark)


**Solution:**

*For Example:*

To cause a shift to the left add  $\text{NO}_2$  **or** remove  $\text{N}_2\text{O}_4$  **or** decrease the volume.

← 1 mark

3. Consider the data obtained for the following equilibrium:



	$[\text{Fe}^{3+}]$	$[\text{SCN}^{-}]$	$[\text{FeSCN}^{2+}]$
Experiment 1	$3.91 \times 10^{-2}$	$8.02 \times 10^{-5}$	$9.22 \times 10^{-4}$
Experiment 2	$6.27 \times 10^{-3}$	$3.65 \times 10^{-4}$	?

Calculate the  $[\text{FeSCN}^{2+}]$  in experiment #2.

**(3 marks)**

**Solution:**

$$\begin{aligned}
 K_{eq} &= \frac{[\text{FeSCN}^{2+}]}{[\text{Fe}^{3+}][\text{SCN}^{-}]} \\
 &= \frac{9.22 \times 10^{-4}}{(3.91 \times 10^{-2})(8.02 \times 10^{-5})} \\
 &= 2.94 \times 10^2
 \end{aligned}
 \left. \vphantom{\begin{aligned} K_{eq} &= \frac{[\text{FeSCN}^{2+}]}{[\text{Fe}^{3+}][\text{SCN}^{-}]} \\ &= \frac{9.22 \times 10^{-4}}{(3.91 \times 10^{-2})(8.02 \times 10^{-5})} \\ &= 2.94 \times 10^2 \end{aligned}} \right\} \leftarrow 1\frac{1}{2} \text{ mark}$$


$$\begin{aligned}
 2.94 \times 10^2 &= \frac{x}{(6.27 \times 10^{-3})(3.65 \times 10^{-4})} \\
 [\text{FeSCN}^{2+}] = x &= 6.73 \times 10^{-4} \text{ M}
 \end{aligned}
 \left. \vphantom{\begin{aligned} 2.94 \times 10^2 &= \frac{x}{(6.27 \times 10^{-3})(3.65 \times 10^{-4})} \\ [\text{FeSCN}^{2+}] = x &= 6.73 \times 10^{-4} \text{ M} \end{aligned}} \right\} \leftarrow 1\frac{1}{2} \text{ mark}$$

(Deduct  $\frac{1}{2}$  mark for incorrect significant figures.)

4. At 25°C, will a precipitate form when 25.0 mL of 0.010 M  $\text{Pb}(\text{NO}_3)_2$  is combined with 75.0 mL of 0.010 M NaI? Support your answer with calculations. (3 marks)

**Solution:**

*For Example:*



$$[\text{Pb}^{2+}] = 0.010 \text{ M} \times \frac{25.0 \text{ mL}}{100.0 \text{ mL}} = 0.00250 \text{ M}$$

$$[\text{I}^{-}] = 0.010 \text{ M} \times \frac{75.0 \text{ mL}}{100.0 \text{ mL}} = 0.00750 \text{ M}$$

} ← 1 mark

$$\text{Trial } K_{sp} = [\text{Pb}^{2+}][\text{I}^{-}]^2$$


$$= (0.00250)(0.00750)^2$$

$$= 1.4 \times 10^{-7}$$

} ← 1 mark

Since Trial  $K_{sp}$  ( $1.4 \times 10^{-7}$ ) >  $K_{sp}$  ( $8.5 \times 10^{-9}$ ) a precipitate does form. } ← 1 mark


5. When HCl is added to a saturated solution of  $\text{CuC}_2\text{O}_4$ , some precipitate dissolves. However, when HCl is added to a saturated solution of  $\text{PbCl}_2$ , additional precipitate forms.



Explain these observations. Support your explanation with chemical equations. **(3 marks)**

**Solution:**

**For Example:**



$\text{H}^+$  from the acid reacts with the  $\text{C}_2\text{O}_4^{2-}$  to form  $\text{HC}_2\text{O}_4^-$  reducing the  $[\text{C}_2\text{O}_4^{2-}]$  and causing a shift to the product side. }  $\leftarrow$  **1 mark**



The common ion effect causes a shift to the right. }  $\leftarrow$  **1 mark**

6. A 0.100 M solution of an unknown weak acid, HX, has a pH = 1.414.  
What is the  $K_a$  for HX?

(4 marks)

**Solution:**

*For Example:*

$$[\text{H}_3\text{O}^+] = 0.03855 \text{ M}$$

← 1 mark

	HX	+ H <sub>2</sub> O	$\rightleftharpoons$	H <sub>3</sub> O <sup>+</sup>	+ X <sup>-</sup>	
[I]	0.100			0	0	
[C]	-0.03855			+0.03855	+0.03855	
[E]	0.061			0.03855	0.03855	

← 2 marks

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{X}^-]}{[\text{HX}]}$$

← 1 mark

$$K_a = \frac{(0.03855)(0.03855)}{0.061}$$

$$= 0.024$$

(Deduct  $\frac{1}{2}$  mark for incorrect significant figures.)

7. Consider the salt ammonium acetate,  $\text{NH}_4\text{CH}_3\text{COO}$ .

a) Write the equation for the dissociation of  $\text{NH}_4\text{CH}_3\text{COO}$ .

(1 mark)

**Solution:**




← 1 mark

b) Write equations for the hydrolysis reactions which occur.

(2 marks)

**Solution:**



} ← 2 marks

c) Explain why a solution of  $\text{NH}_4\text{CH}_3\text{COO}$  has a  $\text{pH} = 7.00$ .  
Support your answer with calculations.

(2 marks)

**Solution:**

$$K_a \text{ for } \text{NH}_4^+ = 5.6 \times 10^{-10}$$

$$K_b \text{ for } \text{CH}_3\text{COO}^- = \frac{1.0 \times 10^{-14}}{1.8 \times 10^{-5}} = 5.6 \times 10^{-10}$$

$$\text{the } K_a \text{ for } \text{NH}_4^+ = K_b \text{ for } \text{CH}_3\text{COO}^-$$

∴ the acidic cation is completely neutralized by the basic anion.

} ← 2 marks


8. The metals Rh, Ti, Cr and Pd are individually placed in 1.0 M solutions of  $\text{Rh}^{2+}$ ,  $\text{Ti}^{2+}$ ,  $\text{Cr}^{2+}$  and  $\text{Pd}^{2+}$  and the cell voltages of the spontaneous reactions are determined.

ION METAL	$\text{Rh}^{2+}$	$\text{Ti}^{2+}$	$\text{Pd}^{2+}$	$\text{Cr}^{2+}$
Rh		no reaction	0.35 V	no reaction
Ti	2.23 V		2.58 V	?
Pd	no reaction	no reaction		no reaction
Cr	1.51 V	no reaction	1.86 V	

a) Arrange the metals in order of **increasing** strength as reducing agents. **(2 marks)**

**Solution:**


*For Example:*



b) Determine the cell voltage for Ti in a 1.0 M solution of  $\text{Cr}^{2+}$ . **(2 marks)**

**Solution:**

*For Example:*






10. Draw and label an electrochemical cell using a copper anode and having an  $E^\circ$  value  $> 1.00\text{ V}$ .  
(2 marks)

**Solution:**

*For Example:*



$\frac{1}{2}$  mark for suitable cathode — Au for example.

$\frac{1}{2}$  mark for suitable ions —  $\text{Au}^{3+}$  and  $\text{Cu}^{2+}$  for example.

1 mark for diagram being an electrochemical cell, not an electrolytic cell.

**END OF KEY**